

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER  
19-962/S-013**

**Final Printed Labeling**

RHPM Review of Final Printed Labeling  
NDA 19-962/S-013

Date of Submission: January 11, 2001  
Date of Review: January 19, 2001  
Applicant Name: AstraZeneca  
Product Name: Toprol-XL (metoprolol succinate) 25, 50, 100 and 200 mg Tablets

Evaluation:

This submission provides for final printed labeling, revised in accordance with our July 10, 2000 approvable letter and subsequent negotiations at a meeting held on October 23, 2000 between AstraZeneca and the Agency. Draft labeling that was acceptable to the Agency was submitted on December 20, 2000 by AstraZeneca.

The final printed labeling is exactly like the labeling submitted on December 20, 2000 except that under the, "Clinical Endpoints in the MERIT-HF Study" table, "Sudden Death" and "Death Due to worsening heart failure" are indented under, "Cardiovascular mortality" as we requested in a telephone communication.

There were no other changes from the last approved package insert.

Recommendation:

An approval letter should issue for this supplement as set forth under 21 CFR 314.70 (b) (3) [Any change in labeling].

/S/ 2/6/01  
Zelda McDonald, RHPM

cc: orig. NDA  
HFD-110  
HFD-110/McDonald  
HFD-110/Blount  
HF-2

9010201 for FDA submittal Toprol-XL Retail PI

01/03/01 4:35 pm SZT

Art created and printed at 100%

9010201




Rev 01/01

# **TOPROL-XL®** (metoprolol succinate) extended-release tablets

Tablets: 25 mg, 50 mg, 100 mg, and 200 mg

## DESCRIPTION

Toprol-XL, metoprolol succinate, is a beta<sub>1</sub>-selective (cardioselective) adrenoceptor blocking agent, for oral administration, available as extended release tablets. Toprol-XL has been formulated to provide a controlled and predictable release of metoprolol for once daily administration. The tablets comprise a multiple unit system containing metoprolol succinate in a multitude of controlled release pellets. Each pellet acts as a separate drug delivery unit and is designed to deliver metoprolol continuously over the dosage interval. The tablets contain 23.75, 47.5, 95 and 190 mg of metoprolol succinate equivalent to 25, 50, 100 and 200 mg of metoprolol tartrate, USP, respectively. Its chemical name is (±) 1-(isopropylamino)-3-[p-(2-methoxyethyl)phenoxy]-2-propanol succinate (2:1) (salt). Its structural formula is:



Metoprolol succinate is a white crystalline powder with a molecular weight of 652.8. It is freely soluble in water; soluble in methanol; sparingly soluble in ethanol; slightly soluble in dichloromethane and 2-propanol; practically insoluble in ethyl-acetate, acetone, diethylether and heptane. Inactive ingredients: silicon dioxide, cellulose compounds, sodium stearyl fumarate, polyethylene glycol, titanium dioxide, paraffin.

## CLINICAL PHARMACOLOGY

### General

Metoprolol is a beta<sub>1</sub>-selective (cardioselective) adrenergic receptor blocking agent. This preferential effect is not absolute, however, and at higher plasma concentrations, metoprolol also inhibits beta<sub>2</sub>-adrenoceptors, chiefly located in the bronchial and vascular musculature. Metoprolol has no intrinsic sympathomimetic activity, and membrane-stabilizing activity is detectable only at plasma concentrations much greater than required for beta-blockade. Animal and human experiments indicate that metoprolol slows the sinus rate and decreases AV nodal conduction.

Clinical pharmacology studies have confirmed the beta-blocking activity of metoprolol in man, as shown by (1) reduction in heart rate and cardiac output at rest and upon exercise, (2) reduction of systolic blood pressure upon exercise, (3) inhibition of isoproterenol-induced tachycardia, and (4) reduction of reflex orthostatic tachycardia.

The relative beta<sub>1</sub>-selectivity of metoprolol has been confirmed by the following: (1) In normal subjects, metoprolol is unable to reverse the beta<sub>2</sub>-mediated vasodilating effects of epinephrine. This contrasts with the effect of nonselective beta-blockers, which completely reverse the vasodilating effects of epinephrine. (2) In asthmatic patients, metoprolol reduces FEV<sub>1</sub> and FVC significantly less than a nonselective beta-blocker, propranolol, at equivalent beta<sub>1</sub>-receptor blocking doses.

In five controlled studies in normal healthy subjects, the same daily doses of Toprol-XL and immediate release metoprolol were compared in terms of the extent and duration of beta<sub>1</sub>-blockade produced. Both formulations were given in a dose range equivalent to 100-400 mg of immediate release metoprolol per day. In these studies, Toprol-XL was administered once a day and immediate release metoprolol was administered once to four times a day. A sixth controlled study compared the beta<sub>1</sub>-blocking effects of a 50 mg daily dose of the two formulations. In

reduces FEV<sub>1</sub> and FVC significantly less than a nonselective beta-blocker, propranolol, at equivalent beta<sub>1</sub>-receptor blocking doses.

In five controlled studies in normal healthy subjects, the same daily doses of Toprol-XL and immediate release metoprolol were compared in terms of the extent and duration of beta<sub>1</sub>-blockade produced. Both formulations were given in a dose range equivalent to 100-400 mg of immediate release metoprolol per day. In these studies, Toprol-XL was administered once a day and immediate release metoprolol was administered once to four times a day. A sixth controlled study compared the beta<sub>1</sub>-blocking effects of a 50 mg daily dose of the two formulations. In each study, beta<sub>1</sub>-blockade was expressed as the percent change from baseline in exercise heart rate following standardized submaximal exercise tolerance tests at steady state. Toprol-XL administered once a day, and immediate release metoprolol administered once to four times a day, provided comparable total beta<sub>1</sub>-blockade over 24 hours (area under the beta<sub>1</sub>-blockade versus time curve) in the dose range 100-400 mg. At a dosage of 50 mg once daily, Toprol-XL produced significantly higher total beta<sub>1</sub>-blockade over 24 hours than immediate release metoprolol. For Toprol-XL, the percent reduction in exercise heart rate was relatively stable throughout the entire dosage interval and the level of beta<sub>1</sub>-blockade increased with increasing doses from 50 to 300 mg daily. The effects at peak/trough (i.e. at 24 hours post dosing) were: 14/9, 16/10, 24/14, 27/22 and 27/20% reduction in exercise heart rate for doses of 50, 100, 200, 300 and 400 mg Toprol-XL once a day, respectively. In contrast to Toprol-XL, immediate release metoprolol given at a dose of 50-100 mg once a day, produced a significantly larger peak effect on exercise tachycardia, but the effect was not evident at 24 hours. To match the peak to trough ratio obtained with Toprol-XL over the dosing range of 200 to 400 mg, a t.i.d. to q.i.d. divided dosing regimen was required for immediate release metoprolol. A controlled cross-over study in heart failure patients compared the plasma concentrations and beta<sub>1</sub>-blocking effects of 50 mg immediate release metoprolol administered t.i.d., 100 mg and 200 mg Toprol-XL once daily. A 50 mg dose of immediate release metoprolol t.i.d. produced a peak plasma level of metoprolol similar to the peak level observed with 200 mg of Toprol-XL. A 200 mg dose of Toprol-XL produced a larger effect on suppression of exercised-induced and Holter-monitored heart rate over 24 hours compared to 50 mg t.i.d. of immediate release metoprolol.

The relationship between plasma metoprolol levels and reduction in exercise heart rate is independent of the pharmaceutical formulation. Using the E<sub>max</sub> model, the maximal beta<sub>1</sub>-blocking effect has been estimated to produce a 30% reduction in exercise heart rate. Beta<sub>1</sub>-blocking effects in the range of 30-80% of the maximal effect (corresponding to approximately 8-23% reduction in exercise heart rate) are expected to occur at metoprolol plasma concentrations ranging from 30-540 nmol/L. The concentration-effect curve begins reaching a plateau between 200-300 nmol/L, and higher plasma levels produce little additional beta<sub>1</sub>-blocking effect. The relative beta<sub>1</sub>-selectivity of metoprolol diminishes and blockade of beta<sub>2</sub>-adrenoceptors increases at higher plasma concentrations.

Although beta-adrenergic receptor blockade is useful in the treatment of angina, hypertension, and heart failure there are situations in which sympathetic stimulation is vital. In patients with severely damaged hearts, adequate ventricular function may depend on sympathetic drive. In the presence of AV block, beta-blockade may prevent the necessary facilitating effect of sympathetic activity on conduction. Beta<sub>2</sub>-adrenergic blockade results in passive bronchial constriction by interfering with endogenous adrenergic bronchodilator activity in patients subject to bronchospasm and may also interfere with exogenous bronchodilators in such patients.

In other studies, treatment with Toprol-XL produced an improvement in left ventricular ejection fraction. Toprol-XL was also shown to delay the increase in left ventricular end-systolic and end-diastolic volumes after 6 months of treatment.

#### **Hypertension**

The mechanism of the antihypertensive effects of beta-blocking agents has not been elucidated. However, several possible mechanisms have been proposed: (1) competitive antagonism of catecholamines at peripheral (especially cardiac) adrenergic neuron sites, leading to decreased cardiac output; (2) a central effect leading to reduced sympathetic outflow to the periphery; and (3) suppression of renin activity.

#### **Clinical Trials**

In controlled clinical studies, an immediate release dosage form of metoprolol has been shown to be an effective antihypertensive agent when used alone or as concomitant therapy with thiazide-type diuretics at dosages of 100-450 mg daily. Toprol-XL, in dosages of 100 to 400 mg once daily, has been shown to possess comparable beta<sub>1</sub>-blockade as conventional metoprolol tablets administered two to four times daily. In addition, Toprol-XL administered at a dose of 50 mg once daily has been shown to lower blood pressure 24-hours post-dosing in placebo controlled studies. In controlled, comparative, clinical studies, immediate release metoprolol appeared comparable as an antihypertensive agent to propranolol, methyldopa, and thiazide-type diuretics, and affected both supine and standing blood pressure. Because of variable plasma levels attained with a given dose and lack of a consistent relationship of antihypertensive activity to drug plasma concentration, selection of proper dosage requires individual titration.

#### **Angina Pectoris**

By blocking catecholamine-induced increases in heart rate, in velocity and extent of myocardial contraction, and in blood pressure, metoprolol reduces the oxygen requirements of the heart at any given level of effort, thus making it useful in the long-term management of angina pectoris.

#### **Clinical Trials**

In controlled clinical trials, an immediate release formulation of metoprolol has been shown to be an effective antianginal agent, reducing the number of angina attacks and increasing exercise tolerance. The dosage used in these studies ranged from 100 to 400 mg daily. Toprol-XL, in dosages of 100 to 400 mg once daily, has been shown to possess beta-blockade similar to conventional metoprolol tablets administered two to four times daily.

#### **Heart Failure**

The precise mechanism for the beneficial effects of beta-blockers in heart failure has not been elucidated.

metoprolol has been shown to be an effective antianginal agent, reducing the number of angina attacks and increasing exercise tolerance. The dosage used in these studies ranged from 100 to 400 mg daily. Toprol-XL, in dosages of 100 to 400 mg once daily, has been shown to possess beta-blockade similar to conventional metoprolol tablets administered two to four times daily.

#### Heart Failure

The precise mechanism for the beneficial effects of beta-blockers in heart failure has not been elucidated.

#### Clinical Trials

MERIT-HF was a double-blind, placebo-controlled study of Toprol-XL conducted in 14 countries including the US. It randomized 3991 patients (1990 to Toprol-XL) with ejection fraction  $\leq 0.40$  and NYHA Class II-IV heart failure attributable to ischemia, hypertension, or cardiomyopathy. The protocol excluded patients with contraindications to beta-blocker use, those expected to undergo heart surgery, and those within 28 days of myocardial infarction or unstable angina. The primary endpoints of the trial were (1) all-cause mortality plus all-cause hospitalization (time to first event), and (2) all-cause mortality. Patients were stabilized on optimal concomitant therapy for heart failure, including diuretics, ACE inhibitors, cardiac glycosides, and nitrates. At randomization, 41% of patients were NYHA Class II, 55% NYHA Class III; 65% of patients had heart failure attributed to ischemic heart disease; 44% had a history of hypertension; 25% had diabetes mellitus; 48% had a history of myocardial infarction. Among patients in the trial, 90% were on diuretics, 89% were on ACE inhibitors, 64% were on digitalis, 27% were on a lipid-lowering agent, 37% were on an oral anticoagulant, and the mean ejection fraction was 0.28. The mean duration of follow-up was one year. At the end of the study, the mean daily dose of Toprol-XL was 159 mg.

The trial was terminated early for a statistically significant reduction in all-cause mortality (34%, nominal  $p=0.00009$ ). The risk of all-cause mortality plus all-cause hospitalization was reduced by 19% ( $p=0.00012$ ). The trial also showed improvements in heart failure-related mortality and heart failure-related hospitalizations, and NYHA functional class.


The table below shows the principal results for the overall study population. The figure below illustrates principal results for a wide variety of subgroup comparisons, including US vs. non-US populations (the latter of which was not pre-specified). The combined endpoints of all-cause mortality plus all-cause hospitalization and of mortality plus heart failure hospitalization showed consistent effects in the overall study population and the subgroups, including women and the US population. However, in the US subgroup and women, overall mortality and cardiovascular mortality appeared less affected. Analyses of female and US patients were carried out because they each represented about 25% of the overall population. Nonetheless, subgroup analyses can be difficult to interpret and it is not known whether these represent true differences or chance effects.

Clinical Endpoint	Number of Patients Placebo n=2001	Toprol-XL n=1990	Relative Risk (95% CI)	Risk Reduction w/Toprol-XL	Nominal P-value
All-cause mortality plus all-cause hospitalization†	767	641	0.81 (0.73-0.90)	19%	0.00012
All-cause mortality	217	145	0.66 (0.53-0.81)	34%	0.00009
All-cause mortality plus heart failure hospitalization†	439	311	0.69 (0.60-0.80)	31%	0.000008
Cardiovascular mortality	203	128	0.62 (0.50-0.78)	38%	0.000022
Sudden death	132	79	0.59 (0.45-0.78)	41%	0.0002
Death due to worsening heart failure	58	30	0.51 (0.33-0.79)	49%	0.0023
Hospitalizations due to worsening heart failure†	451	317	N/A	N/A	0.0000076
Cardiovascular hospitalization†	773	649	N/A	N/A	0.00028

† Time to first event

‡ Comparison of treatment groups examines the number of hospitalizations (Wilcoxon test); relative risk and risk reduction are not applicable.

#### Results for Subgroups in MERIT-HF



US = United States; NYHA = New York Heart Association; EF = ejection fraction; MI = myocardial infarction; HR = heart rate.

#### Pharmacokinetics

In man, absorption of metoprolol is rapid and complete. Plasma levels following oral administration of conventional metoprolol tablets, however, approximate 50% of levels following intravenous administration, indicating about 50% first-pass metabolism. Metoprolol crosses the blood-brain barrier and has been reported in the CSF in a concen-

**Pharmacokinetics**

In man, absorption of metoprolol is rapid and complete. Plasma levels following oral administration of conventional metoprolol tablets, however, approximate 50% of levels following intravenous administration, indicating about 50% first-pass metabolism. Metoprolol crosses the blood-brain barrier and has been reported in the CSF in a concentration 78% of the simultaneous plasma concentration.

Plasma levels achieved are highly variable after oral administration. Only a small fraction of the drug (about 12%) is bound to human serum albumin. Metoprolol is a racemic mixture of R- and S- enantiomers, and is primarily metabolized by CYP2D6. When administered orally, it exhibits stereoselective metabolism that is dependent on oxidation phenotype. Elimination is mainly by biotransformation in the liver, and the plasma half-life ranges from approximately 3 to 7 hours. Less than 5% of an oral dose of metoprolol is recovered unchanged in the urine; the rest is excreted by the kidneys as metabolites that appear to have no beta blocking activity. Following intravenous administration of metoprolol, the urinary recovery of unchanged drug is approximately 10%. The systemic availability and half-life of metoprolol in patients with renal failure do not differ to a clinically significant degree from those in normal subjects. Consequently, no reduction in dosage is usually needed in patients with chronic renal failure.

Metoprolol is metabolized predominantly by CYP2D6, an enzyme that is absent in about 8% of Caucasians (poor metabolizers) and about 2% of most other populations. CYP2D6 can be inhibited by a number of drugs. Concomitant use of inhibiting drugs in poor metabolizers will increase blood levels of metoprolol several fold, decreasing metoprolol's cardioselectivity. (See PRECAUTIONS, Drug Interactions.)

In comparison to conventional metoprolol, the plasma metoprolol levels following administration of Toprol-XL are characterized by lower peaks, longer time to peak and significantly lower peak to trough variation. The peak plasma levels following once daily administration of Toprol-XL average one-fourth to one-half the peak plasma levels obtained following a corresponding dose of conventional metoprolol, administered once daily or in divided doses. At steady state the average bioavailability of metoprolol following administration of Toprol-XL, across the dosage range of 50 to 400 mg once daily, was 77% relative to the corresponding single or divided doses of conventional metoprolol. Nevertheless, over the 24 hour dosing interval,  $\beta_1$ -blockade is comparable and dose-related (see CLINICAL PHARMACOLOGY). The bioavailability of metoprolol shows a dose-related, although not directly proportional, increase with dose and is not significantly affected by food following Toprol-XL administration.

**INDICATIONS AND USAGE****Hypertension**

Toprol-XL is indicated for the treatment of hypertension. It may be used alone or in combination with other antihypertensive agents.

**Angina Pectoris**

Toprol-XL is indicated in the long-term treatment of angina pectoris.

**Heart Failure**

Toprol-XL is indicated for the treatment of stable, symptomatic (NYHA Class II or III) heart failure of ischemic, hypertensive, or cardiomyopathic origin. It was studied in patients already receiving ACE inhibitors, diuretics, and, in the majority of cases, digitalis. In this population, Toprol-XL decreased the rate of mortality plus hospitalization, largely through a reduction in cardiovascular mortality and hospitalizations for heart failure.

**CONTRAINDICATIONS**

Toprol-XL is contraindicated in severe bradycardia, heart block greater than first degree, cardiogenic shock, decompensated cardiac failure, and sick sinus syndrome (unless a permanent pacemaker is in place) (see WARNINGS).

**WARNINGS**

**Ischemic Heart Disease:** Following abrupt cessation of therapy with certain beta-blocking agents, exacerbations of angina pectoris and, in some cases, myocardial infarction have occurred. When discontinuing chronically administered Toprol-XL, particularly in patients with ischemic heart disease, the dosage should be gradually reduced over a period of 1-2 weeks and the patient should be carefully monitored. If angina markedly worsens or acute coronary insufficiency develops, Toprol-XL administration should be reinstated promptly, at least temporarily, and other measures appropriate for the management of unstable angina should be taken. Patients should be warned against interruption or discontinuation of therapy without the physician's advice. Because coronary artery disease is common and may be unrecognized, it may be prudent not to discontinue Toprol-XL therapy abruptly even in patients treated only for hypertension.

**Bronchospastic Diseases:** PATIENTS WITH BRONCHOSPASTIC DISEASES SHOULD, IN GENERAL, NOT RECEIVE BETA-BLOCKERS. Because of its relative  $\beta_1$ -selectivity, however, Toprol-XL may be used with caution in patients with bronchospastic disease who do not respond to, or cannot tolerate, other antihypertensive treatment. Since  $\beta_1$ -selectivity is not absolute, a  $\beta_2$ -stimulating agent should be administered concomitantly, and the lowest possible dose of Toprol-XL should be used (see DOSAGE AND ADMINISTRATION).

**Major Surgery:** The necessity or desirability of withdrawing beta-blocking therapy prior to major surgery is controversial; the impaired

(CONTINUED ON REVERSE SIDE)

Artist Job Notes - This information is not reviewed/approved by QA. These notes are for use only by the AstraZeneca Graphics Department. Refer to the approved specification for details regarding this artwork.  
 System: QuarkXPress 4.01  
 iRc: A5B  
 Fonts: Helvetica 55 Roman, 56 Italic, 75 Bold, 76 Bold Italic; Cheltenham Book, Bold  
 Dimensions: 36-3/4" x 2-3/4"  
 Barcode: 8 digit 12 of 5  
 Color: Black

## TOPROL-XL® (metoprolol succinate) Tablets

ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anesthesia and surgical procedures.

Toprol-XL like other beta-blockers, is a competitive inhibitor of beta-receptor agonists, and its effects can be reversed by administration of such agents, e.g., dobutamine or isoproterenol. However, such patients may be subject to protracted severe hypotension. Difficulty in restarting and maintaining the heart beat has also been reported with beta-blockers.

**Diabetes and Hypoglycemia:** Toprol-XL should be used with caution in diabetic patients if a beta-blocking agent is required. Beta-blockers may mask tachycardia occurring with hypoglycemia, but other manifestations such as dizziness and sweating may not be significantly affected.

**Thyrototoxicosis:** Beta-adrenergic blockade may mask certain clinical signs (e.g., tachycardia) of hyperthyroidism. Patients suspected of developing thyrotoxicosis should be managed carefully to avoid abrupt withdrawal of beta-blockade, which might precipitate a thyroid storm.

### PRECAUTIONS

#### General

Toprol-XL should be used with caution in patients with impaired hepatic function.

Worsening cardiac failure may occur during up-titration of Toprol-XL. If such symptoms occur, diuretics should be increased and the dose of Toprol-XL should not be advanced until clinical stability is restored (see DOSAGE AND ADMINISTRATION). It may be necessary to lower the dose of Toprol-XL or temporarily discontinue it. Such episodes do not preclude subsequent successful titration of Toprol-XL.

#### Information for Patients

Patients should be advised to take Toprol-XL regularly and continuously, as directed, preferably with or immediately following meals. If a dose should be missed, the patient should take only the next scheduled dose (without doubling it). Patients should not interrupt or discontinue Toprol-XL without consulting the physician.

Patients should be advised (1) to avoid operating automobiles and machinery or engaging in other tasks requiring alertness until the patient's response to therapy with Toprol-XL has been determined; (2) to contact the physician if any difficulty in breathing occurs; (3) to inform the physician or dentist before any type of surgery that he or she is taking Toprol-XL.

Heart failure patients should be advised to consult their physician if they experience signs or symptoms of worsening heart failure such as weight gain or increasing shortness of breath.

#### Laboratory Tests

Clinical laboratory findings may include elevated levels of serum transaminase, alkaline phosphatase, and lactate dehydrogenase.

#### Drug Interactions

Catecholamine-depleting drugs (e.g., reserpine) may have an additive effect when given with beta-blocking agents. Patients treated with Toprol-XL plus a catecholamine depletor should therefore be closely observed for evidence of hypotension or marked bradycardia, which may produce vertigo, syncope, or postural hypotension.

Drugs that inhibit CYP2D6 such as quinidine, fluoxetine, paroxetine, and propafenone are likely to increase metoprolol concentration. In healthy subjects with CYP2D6 extensive metabolizer phenotype, coadministration of quinidine 100 mg and immediate release metoprolol 200 mg, tripled the concentration of S-metoprolol and doubled the metoprolol elimination half-life. In four patients with cardiovascular disease, coadministration of propafenone 150 mg t.i.d. with immediate release metoprolol 50 mg t.i.d. resulted in two- to five-fold increases in the steady-state concentration of metoprolol. These increases in plasma concentration would decrease the cardioselectivity of metoprolol.

#### Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have been conducted to evaluate the carcinogenic potential of metoprolol tartrate. In 2-year studies in rats at three oral dosage levels of up to 800 mg/kg/day (41 times, on a mg/m<sup>2</sup> basis, the daily dose of 200 mg for a 60-kg patient), there was no increase in the development of spontaneously occurring benign or malignant neoplasms of any type. The only histologic changes that appeared to be drug related were an increased incidence of generally mild focal accumulation of foamy macrophages in pulmonary alveoli and a slight increase in biliary hyperplasia. In a 21-month study in Swiss albino mice at three oral dosage levels of up to 750 mg/kg/day (18 times, on a mg/m<sup>2</sup> basis, the daily dose of 200 mg for a 60-kg patient), benign lung tumors (small adenomas) occurred more frequently in female mice receiving the highest dose than in untreated control animals. There was no increase in malignant or total (benign plus malignant) lung tumors, nor in the overall incidence of tumors or malignant tumors. This 21-month study was repeated in CD-1 mice, and no statistically or biologically significant differences were observed between treated and control mice of either sex for any type of tumor.

All genotoxicity tests performed on metoprolol tartrate (a dominant lethal study in mice, chromosome studies in somatic cells, a Salmonella/mammalian-microsome mutagenicity test, and a nucleus anomaly test in somatic interphase nuclei) and metoprolol succinate (a Salmonella/mammalian-microsome mutagenicity test) were negative.

No evidence of impaired fertility due to metoprolol tartrate was observed in a study performed in rats at doses up to 22 times, on a mg/m<sup>2</sup> basis, the daily dose of 200 mg in a 60-kg patient.

#### Pregnancy Category C

Metoprolol tartrate has been shown to increase post-implantation loss and decrease neonatal survival in rats at doses up to 22 times, on a

Salmonella/mammalian-microsome mutagenicity test, and a nucleus anomaly test in somatic interphase nuclei) and metoprolol succinate (a Salmonella/mammalian-microsome mutagenicity test) were negative.

No evidence of impaired fertility due to metoprolol tartrate was observed in a study performed in rats at doses up to 22 times, on a mg/m<sup>2</sup> basis, the daily dose of 200 mg in a 60-kg patient.

#### **Pregnancy Category C**

Metoprolol tartrate has been shown to increase post-implantation loss and decrease neonatal survival in rats at doses up to 22 times, on a mg/m<sup>2</sup> basis, the daily dose of 200 mg in a 60-kg patient. Distribution studies in mice confirm exposure of the fetus when metoprolol tartrate is administered to the pregnant animal. These studies have revealed no evidence of impaired fertility or teratogenicity. There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

#### **Nursing Mothers**

Metoprolol is excreted in breast milk in very small quantities. An infant consuming 1 liter of breast milk daily would receive a dose of less than 1 mg of the drug. Caution should be exercised when Toprol-XL is administered to a nursing woman.

#### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

#### **Geriatric Use**

Clinical studies of Toprol-XL in hypertension did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience in hypertensive patients has not identified differences in responses between elderly and younger patients.

Of the 1,990 patients with heart failure randomized to Toprol-XL in the MERIT-HF trial, 50% (990) were 65 years of age and older and 12% (238) were 75 years of age and older. There were no notable differences in efficacy or the rate of adverse events between older and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

#### **Risk of Anaphylactic Reactions**

While taking beta-blockers, patients with a history of severe anaphylactic reactions to a variety of allergens may be more reactive to repeated challenge, either accidental, diagnostic or therapeutic. Such patients may be unresponsive to the usual doses of epinephrine used to treat allergic reaction.

### **ADVERSE REACTIONS**

#### **Hypertension and Angina**

Most adverse effects have been mild and transient. The following adverse reactions have been reported for metoprolol tartrate.

**Central Nervous System:** Tiredness and dizziness have occurred in about 10 of 100 patients. Depression has been reported in about 5 of 100 patients. Mental confusion and short-term memory loss have been reported. Headache, somnolence, nightmares, and insomnia have also been reported.

**Cardiovascular:** Shortness of breath and bradycardia have occurred in approximately 3 of 100 patients. Cold extremities; arterial insufficiency, usually of the Raynaud type; palpitations; congestive heart failure; peripheral edema; syncope; chest pain; and hypotension have been reported in about 1 of 100 patients (see CONTRAINDICATIONS, WARNINGS and PRECAUTIONS).

**Respiratory:** Wheezing (bronchospasm) and dyspnea have been reported in about 1 of 100 patients (see WARNINGS).

**Gastrointestinal:** Diarrhea has occurred in about 5 of 100 patients. Nausea, dry mouth, gastric pain, constipation, flatulence, digestive tract disorders and heartburn have been reported in about 1 of 100 patients.

**Hypersensitive Reactions:** Pruritus or rash have occurred in about 5 of 100 patients. Worsening of psoriasis has also been reported.

**Miscellaneous:** Peyronie's disease has been reported in fewer than 1 of 100,000 patients. Musculoskeletal pain, blurred vision, decreased libido and tinnitus have also been reported.

There have been rare reports of reversible alopecia, agranulocytosis, and dry eyes. Discontinuation of the drug should be considered if any such reaction is not otherwise explicable. The oculomucocutaneous syndrome associated with the beta-blocker practolol has not been reported with metoprolol.

#### **Potential Adverse Reactions**

A variety of adverse reactions not listed above have been reported with other beta-adrenergic blocking agents and should be considered potential adverse reactions to Toprol-XL.

**Central Nervous System:** Reversible mental depression progressing to cataplexy; an acute reversible syndrome characterized by disorientation for time and place, short-term memory loss, emotional lability, slightly clouded sensorium, and decreased performance on neuropsychometrics.

**Cardiovascular:** Intensification of AV block (see CONTRAINDICATIONS).

**Hematologic:** Agranulocytosis, nonthrombocytopenic purpura, thrombocytopenic purpura.

**Hypersensitive Reactions:** Fever combined with aching and sore throat, laryngospasm, and respiratory distress.

#### **Heart Failure**

In the MERIT-HF study, serious adverse events and adverse events leading to discontinuation of study medication were systematically collected. In the MERIT-HF study comparing Toprol-XL in daily doses up to 200 mg (mean dose 159 mg once-daily) (n=1990) to placebo (n=2001), 10.3% of Toprol-XL patients discontinued for adverse events vs. 12.2% of placebo patients.

The table below lists adverse events in the MERIT-HF study that occurred at an incidence of equal to or greater than 1% in the Toprol-XL group and greater than placebo by more than 0.5%, regardless of the assessment of causality.

**Adverse Events Occurring in the MERIT-HF Study at an Incidence  $\geq$  1% in the Toprol-XL Group and Greater than Placebo by More than 0.5%**

T09mLV1      D09m0h



dose 159 mg once-daily) (n=1990) to placebo (n=2001), 10.3% of Toprol-XL patients discontinued for adverse events vs. 12.2% of placebo patients. The table below lists adverse events in the MERIT-HF study that occurred at an incidence of equal to or greater than 1% in the Toprol-XL group and greater than placebo by more than 0.5%, regardless of the assessment of causality.

**Adverse Events Occurring in the MERIT-HF Study at an Incidence  $\geq$  1% in the Toprol-XL Group and Greater than Placebo by More Than 0.5%**

	Toprol-XL n=1990	Placebo n=2001
	% of patients	% of patients
Dizziness/vertigo	1.8	1.0
Bradycardia	1.5	0.4
Accident and/or injury	1.4	0.8

Other adverse events with an incidence of  $> 1\%$  on Toprol-XL and as common on placebo (within 0.5%) included myocardial infarction, pneumonia, cerebrovascular disorder, chest pain, dyspnea/dyspnea aggravated, syncope, coronary artery disorder, ventricular tachycardia/arrhythmia aggravated, hypotension, diabetes mellitus/diabetes mellitus aggravated, abdominal pain, and fatigue.

## OVERDOSAGE

### Acute Toxicity

There have been a few reports of overdosage with Toprol-XL and no specific overdosage information was obtained with this drug, with the exception of animal toxicology data. However, since Toprol-XL (metoprolol succinate salt) contains the same active moiety, metoprolol, as conventional metoprolol tablets (metoprolol tartrate salt), the recommendations on overdosage for metoprolol conventional tablets are applicable to Toprol-XL.

### Signs and Symptoms

Overdosage of Toprol-XL may lead to severe hypotension, sinus bradycardia, atrioventricular block, heart failure, cardiogenic shock, cardiac arrest, bronchospasm, impairment of consciousness/coma, nausea, vomiting, and cyanosis.

### Treatment

In general, patients with acute or recent myocardial infarction or congestive heart failure may be more hemodynamically unstable than other patients and should be treated accordingly. When possible the patient should be treated under intensive care conditions. On the basis of the pharmacologic actions of metoprolol, the following general measures should be employed.

**Elimination of the Drug:** Gastric lavage should be performed.

**Bradycardia:** Atropine should be administered. If there is no response to vagal blockade, isoproterenol should be administered cautiously.

**Hypotension:** A vasopressor should be administered, e.g., levarterenol or dopamine.

**Bronchospasm:** A  $\beta_2$ -stimulating agent and/or a theophylline derivative should be administered.

**Cardiac Failure:** A digitalis glycoside and diuretics should be administered. In shock resulting from inadequate cardiac contractility, administration of dobutamine, isoproterenol or glucagon may be considered.

## DOSAGE AND ADMINISTRATION

Toprol-XL is an extended release tablet intended for once-a-day administration. When switching from immediate release metoprolol tablet to Toprol-XL, the same total daily dose of Toprol-XL should be used.

As with immediate release metoprolol, dosages of Toprol-XL should be individualized and titration may be needed in some patients.

Toprol-XL tablets are scored and can be divided; however, the whole or half tablet should be swallowed whole and not chewed or crushed.

### Hypertension

The usual initial dosage is 50 to 100 mg daily in a single dose, whether used alone or added to a diuretic. The dosage may be increased at weekly (or longer) intervals until optimum blood pressure reduction is achieved. In general, the maximum effect of any given dosage level will be apparent after 1 week of therapy. Dosages above 400 mg per day have not been studied.

### Angina Pectoris


The dosage of Toprol-XL should be individualized. The usual initial dosage is 100 mg daily, given in a single dose. The dosage may be gradually increased at weekly intervals until optimum clinical response has been obtained or there is a pronounced slowing of the heart rate. Dosages above 400 mg per day have not been studied. If treatment is to be discontinued, the dosage should be reduced gradually over a period of 1-2 weeks (see WARNINGS).

### Heart Failure

Dosage must be individualized and closely monitored during up-titration. Prior to initiation of Toprol-XL, the dosing of diuretics, ACE inhibitors, and digitalis (if used) should be stabilized. The recommended starting dose of Toprol-XL is 25 mg once daily for two weeks in patients with NYHA Class II heart failure and 12.5 mg once daily in patients with more severe heart failure. The dose should then be doubled every two weeks to the highest dosage level tolerated by the patient or up to 200 mg of Toprol-XL. If transient worsening of heart failure occurs, it may be treated with increased doses of diuretics, and it may also be necessary to lower the dose of Toprol-XL or temporarily discontinue it. The dose of Toprol-XL should not be increased until symptoms of worsening heart failure have been stabilized. Initial difficulty with titration should not preclude later attempts to introduce Toprol-XL. If heart failure patients experience symptomatic bradycardia, the dose of Toprol-XL should be reduced.

## HOW SUPPLIED

Tablets containing metoprolol succinate equivalent to the indicated weight of metoprolol tartrate, USP, are white, biconvex, film-coated, and scored.

Tablet	Shape	Engraving	Bottle of 100 NDC 0186-
25 mg*	Oval		1088-05

Tablets containing metoprolol succinate equivalent to the indicated weight of metoprolol tartrate, USP, are white, biconvex, film-coated, and scored.

Tablet	Shape	Engraving	Bottle of 100 NDC 0186-
25 mg*	Oval	A B	1088-05
50 mg	Round	A mo	1090-05
100 mg	Round	A ms	1092-05
200 mg	Oval	A my	1094-05

\* The 25 mg tablet is scored on both sides.

Store at 25°C (77°F). Excursions permitted to 15-30°C (59-86°F). (See USP Controlled Room Temperature.)

All trademarks are the property of the AstraZeneca group  
© AstraZeneca 2001

Manufactured for: AstraZeneca LP  
Wilmington, DE 19850  
By: AstraZeneca AB  
S-151 85 Sodertälje, Sweden

Made in Sweden

9010201  
Rev 01/01

AstraZeneca 

